



Jae, S. Y., Kurl, S., Kunutsor, S. K., Heffernan, K., Park, J., & Laukkanen, J. A. (2020). Association between pulse pressure and the risk of sudden cardiac death in middle-aged men: A 26-year follow-up population-based study. *Mayo Clinic Proceedings*.
<https://doi.org/10.1016/j.mayocp.2020.01.033>

Peer reviewed version

License (if available):
CC BY-NC-ND

Link to published version (if available):
[10.1016/j.mayocp.2020.01.033](https://doi.org/10.1016/j.mayocp.2020.01.033)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at [dx.doi.org/10.1016/j.mayocp.2020.01.033](https://doi.org/10.1016/j.mayocp.2020.01.033). Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Research Letter

Association between pulse pressure and the risk of sudden cardiac death in middle-aged men: A 22-year follow-up population-based study

See Young Jae¹, Sudhir Kuri², Setor K. Kunutsor^{3,4}, Kevin Heffernan⁵, Jeong Bae Park⁶,
Jari A. Laukkanen^{2,7,8}

¹Department of Sport Science, University of Seoul, Seoul, Republic of Korea;

²Department of Medicine, Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland;

³National Institute for Health Research Bristol Biomedical Research Centre, University Hospitals Bristol NHS Foundation Trust and University of Bristol, Bristol, UK;

⁴Musculoskeletal Research Unit, Translational Health Sciences, Bristol Medical School, University of Bristol, Learning & Research Building (Level 1), Southmead Hospital, Bristol, BS10 5NB, UK

⁵Department of Exercise Science, Syracuse University, Syracuse, USA

⁶JB Lab and Clinic, Seoul, Republic of Korea

⁷Faculty of Sport and Health Sciences, University of Jyväskylä, Jyväskylä, Finland;

⁸Department of Medicine, Central Finland Health Care District, Jyväskylä, Finland

Corresponding Author: Sae Young Jae, PhD.

Health and Integrative Physiology Laboratory, Department of Sport Science, University of

Seoul. 90 Jeonnong-dong, Dongdaemun-gu, Seoul 130-743, South Korea. E-mail :
syjae@uos.ac.kr, Phone : 82-2-6490-2953 Fax: 82-2-6490-5204

Introduction

Sudden cardiac death (SCD) affects approximately 1 in every 7 adults in the United States.¹ A similarly high burden is reported globally with estimates of 4-5 million cases of SCD per year.² Hypertension increases the risk for SCD, suggesting that blood pressure (BP) is an important risk factor for SCD.³ Pulse pressure (PP) is defined as the numeric difference between systolic and diastolic BP. With advancing age, there are increases in systolic BP with slight decreases in diastolic BP resulting in a widening of PP and associated negligible changes in mean arterial pressure (MAP). Increases in PP are attributable to large artery stiffening and have been shown to be more predictive of adverse cardiovascular outcomes than other BP components, particularly in middle-aged and older adults.^{4, 5} It remains unclear whether PP predicts the risk SCD in the general population. The purpose of this study was to investigate whether PP is associated with the risk of SCD, independently of MAP in middle-aged men.

Methods

Participants were part of the Kuopio Ischaemic Heart Disease Risk Factor Study, which is a prospective population-based study designed to investigate risk factors for CVD and related outcomes in randomly selected sample of men from eastern Finland. At baseline, examinations were conducted on 2,682 men (82.9% of the potential eligible) who resided in the town of Kuopio or its surrounding rural communities between March 1984 and December 1989. Complete data was available on 2,356 participants (aged 42-60) at baseline in this analysis.

Commented [SK1]: This reference is too old. Use this
PMID:
24881994

Resting systolic and diastolic BP were measured using a sphygmomanometer and was expressed as the mean value from 6 different measurements (3 while supine, 1 while standing, and 2 while sitting). Brachial PP was defined as the difference between systolic and diastolic BP. The participants were divided into PP quartiles as follows; quartile 1: <37 mmHg, quartile 2: 37-43 mmHg, quartile 3: 43-51 mmHg, and quartile 4: >51 mmHg.

SCD was defined as a fatal event that occurred within 1 h after the onset of symptoms or within 24 h when autopsy data did not reveal a non-cardiac cause of SCD or after a fatal cardiac arrest following successful resuscitation from ventricular tachycardia and/or ventricular fibrillation. Diagnostic classification of events was based on symptoms, electrocardiographic (ECG) findings, cardiac enzyme elevations, autopsy findings (80% of the SCDs), and history of coronary heart disease (CHD) combined with relevant clinical and ECG findings. Data on SCDs were derived from interviews with family members, hospital records, death certificates, autopsy reports and medico-legal documents.

We used Cox proportional hazard models adjusted for age, body mass index (BMI), high-density lipoprotein cholesterol (HDL-C), low-density lipoprotein cholesterol (LDL-C), triglycerides (TG), C-reactive protein (CRP), diabetes mellitus (DM), hypertension (HTN), family history of CHD, previous myocardial infarction (MI), physical activity, and MAP to determine the hazard ratios (HRs) and 95% confidence intervals (CIs) of PP for SCD. Statistical analyses were conducted using SPSS version 21.0 (SPSS, Armonk, NY).

Results

A total of 253 (10.7%) SCDs occurred during an average of 22 years of follow-up. Cumulative hazard curves demonstrated higher survival among males in the lowest quartile of PP levels compared to those in the highest quartile ($P=xxx$ for log-rank test; **Figure 1**). The HR of SCD in men within the highest quartile of PP ($>51\text{mmHg}$) versus men with the lowest quartile of PP ($<37\text{mmHg}$) was 1.80 (95% confidence interval, 1.18-2.76) after multivariable adjustment. Each unit increment in PP was associated with a 2% (HR, 1.02, 95% CI, 1.01-1.03) increased risk of SCD following adjustment for potential confounders (**Table 1**).

Commented [SK2]: A median (interquartile range) follow-up of xx (xx) years

Discussion

These results suggest that increased PP is associated with a higher risk of SCD in the general population and this is independent of MAP as well as several established and emerging risk factors. The widening of PP with advancing age and CVD is mechanistically linked to large artery stiffness. Increased large artery stiffness detrimentally alters ventricular-vascular coupling and cardiac energetics such that afterload is increased (i.e. increased oxygen demand) and myocardial perfusion reduced (i.e. reduced oxygen supply). Subendocardial ischemia may serve as the substrate for lethal arrhythmogenesis. Large artery stiffness may also hasten the atherosclerotic process, contributing to plaque rupture and myocardial infarction. Atherosclerotic CVD is a notable cause of SCD and responsible for ~50-80% of all cases.⁶

Commented [SK3]: You will need to add a reference somewhere.

Diagnosis, treatment and management of CVD and hypertension-related target organ

damage relies exclusively on the assessment of extreme BP values during the cardiac cycle; the peak (i.e. systolic) and trough (i.e. diastolic). Examination of the BP curve with attention to the pulsatile component of BP (i.e. PP) offers insight on CVD risk extending beyond the assessment of the steady component of the curve (i.e. MAP) or individual extreme values (systolic or diastolic BP). Elevated PP has been shown to be associated with increased risk of incident atrial fibrillation,⁷ heart failure, coronary artery disease and cardiovascular mortality.⁸ Our novel findings add to the literature on the role of PP as an indicator of heart health. Further studies are required to replicate these findings and evaluate if brachial PP may help to identify patients who are at increased risk for SCD and therefore aid in early therapeutic decision-making.


Table 1. Hazard ratio (HR) and 95% confidence interval (CI) for risk of sudden cardiac death by pulse pressure quartiles.

Variables	Case/n (% incident)	Unadjusted HR (95% CI)	Model 1 HR (95% CI)	Model 2 HR (95% CI)
<37 mmHg	41/567 (7.2%)	1 (ref)	1 (ref)	1 (ref)
37-43 mmHg	50/579 (8.6%)	1.24 (0.82-1.87)	1.16 (0.75-1.78)	1.11 (0.72-1.71)
43-51 mmHg	75/628 (11.9%)	1.77 (1.21-2.59)	1.85 (1.25-2.74)	1.67 (1.12-2.50)
>51 mmHg	87/582 (14.9%)	2.43 (1.68-3.53)	2.17 (1.47-3.22)	1.80 (1.18-2.76)
Each 1 mmHg	253/2,356 (10.7%)	1.03 (1.02-1.04)	1.02 (1.01-1.03)	1.02 (1.01-1.03)

Model 1: Adjusted for age, body mass index, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglycerides, C-reactive protein, history of diabetes mellitus, history of hypertension, family history of coronary heart disease, previous myocardial infarction, and physical activity

Model 2: model 1 plus mean arterial pressure

Figure 1. The age and examination year adjusted survival curves of sudden cardiac death according to quartiles of pulse pressure.



References

1. Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, Chamberlain AM, Chang AR, Cheng S, Das SR, Delling FN, Djousse L, Elkind MSV, Ferguson JF, Fornage M, Jordan LC, Khan SS, Kissela BM, Knutson KL, Kwan TW, Lackland DT, Lewis TT, Lichtman JH, Longenecker CT, Loop MS, Lutsey PL, Martin SS, Matsushita K, Moran AE, Mussolino ME, O'Flaherty M, Pandey A, Perak AM, Rosamond WD, Roth GA, Sampson UKA, Satou GM, Schroeder EB, Shah SH, Spartano NL, Stokes A, Tirschwell DL, Tsao CW, Turakhia MP, VanWagner LB, Wilkins JT, Wong SS and Virani SS. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. *Circulation*. 2019;139:e56-e528.
2. Chugh SS, Reinier K, Teodorescu C, Evanado A, Kehr E, Al Samara M, Mariani R, Gunson K and Jui J. Epidemiology of sudden cardiac death: clinical and research implications. *Progress in cardiovascular diseases*. 2008;51:213-28.
3. Verdecchia P, Angeli F, Cavallini C, Aita A, Turturiello D, De Fano M and Reboldi G. Sudden Cardiac Death in Hypertensive Patients. *Hypertension (Dallas, Tex : 1979)*. 2019;73:1071-1078.
4. Franklin SS, Larson MG, Khan SA, Wong ND, Leip EP, Kannel WB and Levy D. Does the relation of blood pressure to coronary heart disease risk change with aging? The Framingham Heart Study. *Circulation*. 2001;103:1245-9.
5. Glynn RJ, Chae CU, Guralnik JM, Taylor JO and Hennekens CH. Pulse pressure and mortality in older people. *Archives of internal medicine*. 2000;160:2765-72.
6. Myerburg RJ and Junttila MJ. Sudden cardiac death caused by coronary heart disease. *Circulation*. 2012;125:1043-52.
7. Mitchell GF, Vasan RS, Keyes MJ, Parise H, Wang TJ, Larson MG, D'Agostino RB, Sr., Kannel WB, Levy D and Benjamin EJ. Pulse pressure and risk of new-onset atrial fibrillation. *Jama*. 2007;297:709-15.
8. Vaccarino V, Holford TR and Krumholz HM. Pulse pressure and risk for myocardial infarction and heart failure in the elderly. *Journal of the American College of Cardiology*. 2000;36:130-8.